



The microbiome of biogas reactors treating lignocellulosic substrates revealed different mechanisms for carbohydrates utilization

Kougias, Panagiotis; Campanaro, S.; Treu, Laura; Tsapekos, Panagiotis; Angelidaki, Irini

Publication date:
2016

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Kougias, P., Campanaro, S., Treu, L., Tsapekos, P., & Angelidaki, I. (2016). *The microbiome of biogas reactors treating lignocellulosic substrates revealed different mechanisms for carbohydrates utilization*. Abstract from 16th International Symposium on Microbial Ecology, Montreal, Canada.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

The microbiome of biogas reactors treating lignocellulosic substrates revealed different mechanisms for carbohydrates utilization

P.G. Kougias¹, S. Campanaro², L. Treu¹, P. Tsapekos¹ and I. Angelidaki¹

¹Department of Environmental Engineering, Technical University of Denmark, Kgs. Lyngby, DK-2800, Denmark

² Department of Biology, University of Padova, Via U. Bassi 58/b, Padova 35121, Italy

Abstract

The present study dissected the microbiome of biogas reactors treating lignocellulosic substrate and swine manure by means of high throughput Illumina sequencing. A comparative metagenomic analysis allowed to identify the microbial species firmly attached to the digested lignocellulosic particles and to distinguish them from the planktonic microbes floating in the liquid medium. *Proteobacteria* and *Firmicutes* were the most abundant phyla identified respectively in the liquid samples and firmly attached to the grass, and accounted approximately 17 and 22% of the total microbial counts. Additionally, *Actinobacteria* were also presented in both samples but in lower relative abundance. Assembly of the shotgun reads followed by a binning process led to the extraction of 151 genome bins, out of which 80 microbial species were completely new and not previously deposited in any database. Moreover, it was shown that 25 microbial genomes were more enriched (>2 fold) in the firmly attached grass samples compared to the liquid phase. A bioinformatic approach based on multiple databases for functional annotation (KEGG, COG, SEED and dbCAN) demonstrated that these microbial species encode enzymes related to carbohydrate utilisation and present numerous carbohydrate binding modules. Finally, it was found that apart from the cellulosome multi-enzyme complex, specific microbes, such as *Bacteroidetes*, present different mechanisms for binding and degrading the lignocellulose due to the presence of multiple CBM6 modules in beta-xylosidase and endoglucanase proteins or SLH modules in unknown proteins.